

# **Areal interpolation of population counts using pre-classified land cover data**

**Population Association of America annual meeting, 2006**

Michael Reibel  
Department of Geography and Anthropology  
California State Polytechnic University, Pomona  
mreibel@csupomona.edu

Aditya Agrawal  
Center for GIS Research  
California State Polytechnic University, Pomona

Current Address:  
Redlands Institute, University of Redlands  
aditya\_agrawal@redlands.edu

**Note: This research is a work in progress. A slightly different version of this paper is under review at *Population Research and Policy Review*.**

## **Abstract**

The need to combine spatial data representing socio-demographic information across various and incompatible spatial units is a common problem for demographers. A particular concern is computing small area trends when aggregation zone boundaries change during the trend interval. To combine population counts across incompatible tract geographies corresponding to successive census enumerations, the authors propose dasymetric areal interpolation using a pre-classified urban land cover data layer. This solution is preferable to traditional re-aggregation techniques. A test of the interpolation technique using the National Land Cover Dataset (NLCD) shows significant error reduction over area weighted interpolation of the same data. The NLCD compares favorably with other common techniques when considered on the basis of accuracy, precision and ease of use.

**Keywords:** areal interpolation, population estimates, census geography, GIS

## **Introduction**

This study proposes accurate areal interpolation as a solution to the common spatial analysis problem of combining incompatible spatial data, e.g. temporal analyses of socio-demographic trends from census tracts. We begin by discussing the problem and the traditional solution of re-aggregation to create a synthetic system of compatible zones. A discussion of areal interpolation using Geographic Information Systems (GIS) follows, including a description of the steps involved that can be followed by demographers with access to a GIS. The authors propose the US Geological Survey's National Land Cover Dataset (NLCD) data set as a highly suitable spatial data layer for performing areal interpolation of spatially aggregated incompatible data.

The authors will then present a case study test of the accuracy of tract level population estimates computed using the NLCD data and dasymetric areal interpolation. Interpolated estimates are benchmarked against a reference set of tract population counts generated from block counts, and the computed errors are mapped, analyzed and compared to the corresponding error distribution of a set of estimates interpolated using a simpler technique with more restrictive assumptions. In light of these findings dasymetric areal interpolation using the NLCD data layer is evaluated relative to alternative techniques with respect to count accuracy, geographic precision, and ease of computation.


## **Theoretical Background**

An important methodological problem in spatial demography is the frequent need to combine incompatible spatial data. Incompatibility, in such situations, means data must be combined in a single record for analysis that are aggregated (or coded, in the case of microdata) to two or more superimposed zone systems, and that in some local areas within the region the zone boundaries of one zone system cross the boundaries of the other zone system. If boundaries of zones in the study area never cross boundaries of the other, superimposed zone system, the systems are nested with respect to each other. In nested zone systems, zones are either a perfect match in each system or, in other cases, whole zones in one zone system can be reaggregated to areas corresponding perfectly to one or more zones in the other system and the counts summed.

The typical problem of incompatible spatial data in demography is the need to combine tract level population and subpopulation count data for the same region pertaining to two successive census enumerations, in order to compute an exhaustive and mutually exclusive set of tract level trends for the time interval. Other studies, notably in the contextual effects modeling of microdata and in the metropolitan analysis of microdata, have encountered similar difficulty in the geoprocessing of data aggregated to incompatible area units.

In the past, investigators have typically sought to solve the problem of combining spatially mismatched data by re-aggregating area units from two superimposed zone systems to the zone coverage containing the smallest possible compatible re-aggregation zones. Compatible re-aggregation zones are a synthetic zone system consisting only of one or more whole area units in both original zone systems to which data are aggregated. It is not always necessary for zones to be nested in order to create compatible re-aggregation zones. It is possible to re-aggregate areas where zone boundaries from one system cross the boundaries of the second system's zones using only whole zones from both zone systems. But this normally requires that a larger number of zones in both systems be combined, often resulting in synthetic re-aggregation zones that are much bigger than the normal scale of the original zones. In Figure 1 we can see certain areas, for

example in the south and southeast of the study area, that experienced rapid population growth between 1990 and 2000. Consequently, many of the 1990 tracts in these areas split into two, three or more new tracts. Moreover, in some cases, these fragments of former tracts were re-aggregated into new tracts that combine territory from two or (rarely) more previous tracts in order to maintain near-target tract population levels. In other words, many 1990 tracts did not neatly split or merge with other 1990 tracts when the 2000 tract geography was created, but rather in a number of cases a 1990 tract was simultaneously split and partially merged.



*Figure 1. Temporal change across complex zone geographies (US Census tracts)*

Superimposed zone systems from different, non-standardized sources thus routinely cross each other's boundaries. The complex territorial recombinations typical of geographic area changes over time for census tracts and other data series aggregation zones, and other such mismatches, frequently lead to situations in which systems of hundreds of zones would need to be reduced to fewer than ten re-aggregation zones in order to match geographies. Such an outcome renders appropriate-scale analysis of local variation within a region impossible. To cope with these situations, investigators must choose between extreme exaggeration in the scale of some zones included in their analyses, or deleting the observations associated with the troublesome zones.

Areal interpolation is a more elegant and, with the use of geographic information systems (GIS), often easier solution to matching incompatible data aggregation zones than the re-aggregation approaches described above (cf. Eicher and Brewer, 2001; Goodchild et al., 1993; Gotway and Young 2002). Moreover, because re-aggregation inevitably involves situations in

which complex boundary changes leave analysts with no solution that would preserve both the processing protocol and the appropriate scale of area units, areal interpolation can be assumed to be more geographically precise and more reliably exhaustive of the study area territory. Areal interpolation refers broadly to techniques that assign data from one or more sets of geographic areas to which data are aggregated (the source zones) to another incompatible and superimposed set (the target zones) using spatial algorithms. In current practice, most of these algorithms exploit the map overlay capabilities of GIS (Longley et al., 2005).

The simplest type of areal interpolation is area weighted interpolation, which requires no information besides the geography of both sets of zone units and the counts to be interpolated from the source to the target zones (Goodchild and Lam, 1980). In area weighted interpolation the two incompatible zone systems describing a given region are superimposed and intersected, creating a set of intersection zones, each of which describes a unique pair of one source and one target zone (Flowerdew and Green, 1992; Reibel and Bufalino, 2005). Each intersection zone is assigned a fraction of its respective source zone's count corresponding to the proportion of the source zone's area occupied by the intersection zone. The intersection zone counts can then be summed across their respective target zones to complete the integration of data to the incompatible zone system.

It is immediately apparent that area weighted interpolation relies on the assumption that there are no internal variations in count density within any source zone, an assumption that is not generally warranted. All other area interpolation techniques seek to improve the accuracy of estimates by bringing to bear meaningful information regarding local density variations of counts within the source zones. Pycnophylactic smoothing techniques use the density surface of the set of source tract counts themselves, and create fine grained, smooth estimated density gradients inside the source tracts by interpolating each tract's count internally based on the count densities of adjacent tracts (Tobler, 1979). The resulting estimated population surface can be used as locally detailed geographic information as-is, or it can be re-aggregated to another zone system that is incompatible with the first. These smoothing techniques, however, require a relatively high level of geostatistical and geoprocessing skill, and can introduce error when (as is the case with census tracts) count density gradients are not in fact typically smooth up to and beyond tract boundaries. Indeed, census tracts frequently have abrupt population density changes that coincide with the distinctive features in the built environment (such as major highways and strips of abandoned brown fields) that are typically chosen as tract boundaries.

The other general approach to areal interpolation that seeks to apply information about internal source zone density gradients involves dasymetric mapping. Dasymetric maps "depict quantitative areal data using boundaries that divide the mapped area into zones of relative homogeneity with the purpose of best portraying the underlying statistical surface" (Eicher and Brewer, 2001). In the context of areal interpolation, dasymetric techniques mean that a (typically fine grained) ancillary data layer is used as a proxy for count densities. This additional layer is superimposed on the source zones from which counts are to be interpolated, and counts are re-assigned to a higher resolution within the source zones corresponding to the detailed geography of the ancillary data values. The counts are then re-aggregated from this relatively fine grained surface or zone system to the set of target zones.

A considerable variety of ancillary data layers has been brought to bear on the question of dasymetric areal interpolation, with corresponding variation in the difficulty of spatial data processing and the quality of results. Most studies have used remotely sensed urban land cover surface data as a weighting factor, but some have used objects such as the street grid (Reibel and Bufalino, 2005), or control zones corresponding to functional areas (Goodchild et al., 1993; Reibel and Agrawal, 2005). Dasymetric mapping using remotely sensed urban land cover data

(Fisher and Langford 1996, Langford and Unwin 1994) as well as Reibel and Bufalino's street weighting technique, have proven to be relatively accurate. Moreover, the larger number of studies using land cover data, relative to smoothing and other dasymmetric weighting techniques, and the robust tests documenting its accuracy, including re-sampling simulations in Cockings et al., (1997) make land cover weighting the normative approach to areal interpolation. The major difficulty of urban land-cover weighted areal interpolation until now has been the need to transform raw images into an information surface, a difficult procedure called digital image processing or classification (Schowengerdt, 1997; Jensen, 1996; Lillesand and Kiefer, 1987).

## **Data and methods**

This study provides an applied example and test of urban land cover weighted areal interpolation using a detailed pre-classified land cover data layer. These data and methods are a good fit for demographers who use GIS but are not GIS specialists because they offer the power and accuracy of land cover weighted interpolation without the need to classify remotely sensed images. Our approach uses land cover data derived through the National Oceanic and Atmospheric Administration's (NOAA) Coastal Change Analysis (C-CAP)<sup>1</sup> Program. Because these data are integrated into the US Geological Survey's National Land Cover Dataset (NLCD), we hereafter, refer to the dataset as the NLCD. The NLCD data is free, seamless and downloadable from <http://seamless.usgs.gov/website/seamless/viewer.php>. These high quality data, derived from Landsat satellite images, provide pre-classified information on land cover category types, including urban land cover, at 30 meter resolution for the entire United States. The authors will provide a set of steps to guide investigators as they perform interpolations using these data, as well as a discussion and examination of the estimation errors in the interpolated population counts in our example.

In this example, the authors will perform land cover weighted areal interpolation using the NLCD data to interpolate 2000 census tract population counts in a study area in eastern Los Angeles County to the 1990 census tract geography. The study area consists of the San Gabriel Valley region and adjacent areas to the east extending to the border of Los Angeles County (Figure 2). The study area contains cities of over 100,000 persons, heavy industrial areas, dams and spillways, college campuses, old citrus packing towns, low density suburbs and a border of hills and mountains. It is therefore generally representative of the various landscapes within Los Angeles County with the exception of coastal areas and the metropolitan central business district.

---

<sup>1</sup> Information about the NOAA's C-CAP program and its integration into the USGS NLCD effort can be found at <http://www.csc.noaa.gov/crs/lca/ccap.html>.




Figure 2. San Gabriel Valley study area

The interpolation consists of a series of steps, mostly performed in a GIS environment using ArcGIS 9.0 (Environmental Systems Research Institute, Redlands, CA). The weighting regression is performed in the statistical package SPSS (SPSS Inc., Chicago, IL). Most popular GIS packages lack integrated regression and statistical analysis functionalities, but it is a simple matter to transfer tabular data back and forth between the two packages by exporting the tables to Dbase (.dbf) file format, which is accessible by both packages. The first task is to fit a set of raster (grid cell) weights corresponding to inhabited land cover types. To do this, the authors computed the proportion of each source (2000) tract's land area consisting of 30x30 meter grid cells of each land cover type which might reasonably be expected to be inhabited. The tracts' populations were then regressed on the areas of inhabited land cover types using ordinary least squares (OLS) regression using SPSS<sup>2</sup>. The resulting coefficients are the raster population weights for each urban land cover type<sup>3</sup>. The regression derived weights for our best fitting raster land cover model ( $R^2=.873$ ) are given in Table 1.

<sup>2</sup> Flowerdew and Green (1989) note that Poisson regression is theoretically preferable for modeling counts. In this study and in others OLS coefficients tend to be almost identical to Poisson coefficients for the same variables and data in areal interpolation (cf. Fisher and Langford, 1995; Langford et al., 1991). Using OLS gives weights the conceptual simplicity that coefficients are a linear function of population density.

<sup>3</sup> Weighting regressions for areal interpolation should be fitted without intercepts, since areas with no inhabitable land cover are expected to have no population.

Table 1. Weights derived by regression for inhabitable land cover categories

Value	Land Cover Type	Weight
3	High Intensity Urban Residential	0.808
4	Low Intensity Residential	0.374
5	Suburban Residential	0.274

*F* statistic = 809.76 sig = .000


Once the weights have been computed, they are applied to the NLCD data layer's raster surface to generate a population surface map. However, because the model only accounts for 87.3% of the variation in the source zone population, the weighted estimates in the population surface must be scaled by the ratio of their respective source zone's observed population to its fitted population to account for the proportion of source zone population not predicted in the model, thus preserving the pycnophylactic property (Tobler 1979). To do this, the grid cells forming the raw estimated population surface were multiplied by the ratio of their respective source tracts' observed populations to the source tract's fitted population computed by summing the raw estimates across the source tract's grid cells:

$$G_s = G_w \left( \frac{T_G}{\hat{T}_G} \right)$$

Where  $G_s$  is the scaled population estimate of grid cell  $G$ ,  $G_w$  is the raw weighted population estimate of grid cell  $G$ ,  $T_G$  is the observed population of the source tract of grid cell  $G$ , and  $\hat{T}_G$  is the fitted population of the source tract of grid cell  $G$  derived by applying the weights to all grid cells and summing across all grid cells in source tract  $T$ .

The result is the scaled population estimate surface map shown in Figure 3. Figure 3 shows the highly erratic and discontinuous nature of the study area's population distribution. Most of the less populated (thus lightly shaded) areas consist of steep hills; with two major exceptions: the San Gabriel River flood plain and quarrying area running southwest from the north central part of the map, and the major industrial corridor along the Pomona Freeway that shows up as a light colored upturned crescent in the southern central part of the map. Dense urban areas are Pomona in the east, Pasadena in the northwest, and in the southwest, smaller dense concentrations in the blue collar suburbs of Alhambra, Montebello, El Monte and La Puente.






*Figure 3. Scaled 2000 tract population counts derived from using the NLCD as a weighting layer where population estimates are now allocated to each grid cell.*

While population surface maps are simply a processing stage in areal interpolation to combine data across incompatible zone systems, such maps are valuable information in themselves, as they are useful for exploratory data analysis and the detailed display of small area populations and/or their characteristics. Next, the grid cells in the population surface map are re-aggregated to the target zone geography, 1990 census tracts in our case, and the grid cell estimates summed across their respective target tracts to yield estimates of the source counts interpolated to the target zones, i.e. the 2000 populations of the 1990 tracts.

The above steps complete the interpolation. Our test of the accuracy of the NLCD land cover weighted interpolated population estimates requires three additional steps: 1) the computation of a set of benchmark counts corresponding to the true 2000 populations of the 1990 tracts; 2) computation of the estimation errors via subtraction from the benchmark counts; and 3) the exploration and analysis of the error distribution. The benchmark counts for the 2000 populations of the 1990 tracts are computed by aggregating the 2000 census block counts to the 1990 census tract geography, using the technique of block centroid aggregation described in Reibel and Bufalino (2005). The errors in estimation computed by subtraction from these benchmark observed counts are explored and mapped. Finally, the error distribution and root mean square (RMS) errors for the NLCD weighted tract estimates and corresponding interpolated estimates from simple area weighting are statistically analyzed and tested for significant improvement in accuracy.


## Results

Figure 4 illustrates the 2000 weighted and scaled population estimates at the grid cell level aggregated to the 1990 tract geography. While both Figures 3 and 4 display the same estimated population distribution, the resolutions differ markedly, and thus, so too does the user's ability to visualize the variation of the population distribution across space. Hence, the methods described in this paper allow for both the generation of a detailed population surface and the creation of interpolated estimates to combine data drawn from superimposed but incompatible zones. In Figure 4, the 1990 census tracts with the largest estimated 2000 population counts are generally those that grew the fastest during the 1990s. Most of these 1990 tracts were split, often in complex ways that included some merges, after the 2000 enumeration to preserve the desired range of tract populations, so it is natural for such 1990 tracts to have very large populations when we use our technique to effectively reverse the splits. In our example, 1990 tracts with estimated 2000 populations of 13,000 to nearly 30,000 can be seen in mostly upscale new hilltop developments in Walnut, Rowland Heights and Diamond Bar in the southeast and La Verne in the northeast, as well as distinctly less prosperous El Monte and parts of north Pasadena further west.



*Figure 4. 2000 population counts aggregated to the 1990 tract geography based on the population surface illustrated in Figure 3.*

Figure 5 illustrates the distribution of estimation errors in the interpolated 2000 population estimates of 1990 tracts in the study area. Spatial autocorrelation tests reveal that the errors are distributed in a spatially dispersed pattern ( $p \leq 0.01$ ). Despite the lack of clustering implied by this observed negative spatial autocorrelation, a close reading of the error map reveals clear patterns of error associated with specific land uses and socio-demographic areas. The high positive errors are concentrated in the industrial areas of Irwindale, El Monte and the City of Industry and in campus neighborhoods including Cal Poly Pomona and Cal Tech in Pasadena. Presumably, some industrial and academic buildings are coded as apartments in these areas, causing overweighting in the estimation process. In the west of the study area, in San Gabriel and Monterey Park, other isolated high positive error tracts are gentrifying neighborhoods completing their transition to from mixed and heavily Hispanic to majority Chinese populations. In the process, many larger houses are being built and inhabited by smaller households. The result is overestimation of the population based on land cover weighting. In the far southeast of the study area is another high positive error tract in upscale Diamond Bar. This neighborhood is new development but it also has similar large homes and small households, and is also heavily Asian, including many Chinese, Filipinos and Koreans.



*Figure 5. Errors in the aggregated 2000 population estimates (Figure 4) compared to the benchmark dataset*

High negative errors are found in poor, less densely built peripheral neighborhoods that are overwhelmingly Hispanic and heavily populated by more recent immigrants. Severe housing overcrowding in such areas, which include western Pomona and parts of Hacienda Heights, La Puente and Covina, creates much larger actual populations than the weighting scheme estimates given the relatively low density residential built environment.

We can interpret from Table 2 that the error distributions for both the NLCD land cover weighted estimates and the area weighted estimates are reasonably symmetrical with means near zero, and that they correspond to approximate Gaussian normality. Overall, the NLCD weighted distribution has fewer large errors in both the negative and positive direction: The maximum and minimum error values are both greater in the area weighted model and the error values at the extremes (>90<sup>th</sup> and <10<sup>th</sup> percentiles) were also greater in the area weighted model. This was expected; the NLCD estimates should perform better because they bring to bear information about the internal count density variations within the source zones from which counts are re-assigned via interpolation. Error reduction was not uniform at every level of the distribution, however. Surprisingly, a range of positive errors in the NLCD weighted distribution, those between the 60<sup>th</sup> and 80<sup>th</sup> percentiles, are in fact slightly larger than the errors at corresponding percentiles of the area weighted distribution.

*Table 2: Error Distributions*

	<i>NLCC Weighted</i>	<i>Area Weighted</i>
Min	-607.5	-890.1
Max	542.8	866.5
<b>Percentiles</b>		
1	-447.0	-585.9
5	-197.3	-236.4
10	-137.3	-156.7
20	-79.3	-83.3
30	-43.6	-50.0
40	-23.2	-20.7
50	-3.0	-4.4
60	15.9	8.7
70	37.1	35.6
80	70.0	67.7
90	134.0	146.8
95	205.3	282.2
99	373.9	591.6

In order to compare overall error levels for the two distributions, and to test whether the observed error reduction associated with one technique over the other is statistically significant, we must compute an overall error statistic. The simple statistic most frequently used for error

distributions is root mean square (RMS) error. RMS error is essentially the standard deviation of the error distribution:

$$\sqrt{\frac{\sum_{i=1}^m (\hat{P}_i - P_i)^2}{m}}$$

Where  $P_i$  is the benchmark population of zone  $i$ ,  $\hat{P}_i$  is the estimated population of zone  $i$ , and  $m$  is the number of zones in the metropolitan or other study area. Table 3 shows that the RMS for the NLCD weighted estimates is 131.93, while that for the area weighted estimates is 173.45. The degree of error reduction achieved by the NLCD weighting over the area weighting is thus 23.94%. To determine whether this error reduction is statistically significant, the authors performed a difference of proportions test on the RMS errors of the two error distributions. The test statistic was highly significant ( $.001 < \text{probability of inference error} < .005$ ). We can conclude that in our example, the dasymetric areal interpolation using the NLCD pre-classified land cover data as a weighting layer achieves a substantial and statistically significant error reduction over area weighted areal interpolation of the same data and study area.

Table 3: Error Analysis

	Area Weighted	NLCD Weighted
N (obs)	281	281
Variance	30084.9	17462.9
Pooled Standard Error	13.00805	
RMS	173.45	131.93
Error Reduction	23.94%	
T Stat, difference of proportions	3.191871	
Significance, difference in RMS	.001 < prob. Error < .005	

## Discussion

The authors have described the problem of combining spatially incompatible data in demographic research, and briefly described two families of solutions to the problem: re-aggregation and estimation by areal interpolation. On the merits, any properly executed areal interpolation, even the relatively crude area weighting technique, will better preserve the scale and exhaustiveness of the zones used for spatial data being processed, i.e. its geographic precision, than will re-aggregation. This is because when mismatches become complex, re-aggregation techniques force investigators to choose between very large re-aggregation zones and dropping problem areas from the analysis – both solutions that are likely to introduce bias in

the analysis of spatial data sets so processed. Moreover, with the use of GIS, areal interpolation is also much easier than re-aggregation.

This study provides both a test of and a guide for using a relatively accurate dasymetric areal interpolation technique with a weighting data layer that does not require digital image processing to classify urban land cover information. The NLCD weighted estimates described in this study are considerably more accurate than area weighted estimates derived from the same data and geography. We believe that the NLCD data, when used for dasymetric areal interpolation, provide a very good combination of accuracy, preservation of data at appropriate scales, and ease of estimation. We hope that demographers will consider areal interpolation when they are performing local and neighborhood analysis that requires combining incompatible spatial data, and we recommend the NLCD weighting approach described here. We also hope that the availability of GIS to help solve difficult data processing problems will facilitate and help promote demographic research on urban and spatial research questions in demography.

## References

- Cockings, S., Fisher P. and Langford, M. (1997). Parameterization and visualization of the errors in areal interpolation. *Geographical Analysis* 29(4): 314–328.
- Eicher, C. and Brewer, C. (2001). Dasymetric mapping and areal interpolation: implementation and evaluation. *Cartography and Geographic Information Science* 28: 125–138.
- Fisher, P. and Langford, M. (1996). Modeling sensitivity to accuracy in classified imagery. *Professional Geographer* 48: 299–309.
- Fisher, P. and Langford, M. (1995). Modeling the errors in areal interpolation between zonal systems by Monte Carlo simulation. *Environment and Planning A* 27: 211–224.
- Flowerdew, R. and Green, M. (1992). Developments in areal interpolation methods and GIS. *Annals of Regional Science* 26: 67–78.
- Flowerdew, R. and Green, M. (1989). Statistical methods for inference between incompatible zone systems, pp. 239-247, in: M. Goodchild and S. Gopal (eds.), *The Accuracy of Spatial Databases*. London: Taylor and Francis
- Goodchild, M., Anselin, L. and Deichmann, U. (1993). A framework for the aerial interpolation of socioeconomic data. *Environment and Planning A* 25: 383–397.
- Goodchild, M. and Lam, N. (1980). Aerial interpolation: a variant of the traditional spatial problem. *Geo-Processing* 1: 297–312.
- Gotway, C. and Young, L., 2002, Combining incompatible data. *Journal of the American Statistical Association* 97: 632–648.
- Jensen, J. R. (1996). *Introductory Digital Image Processing – A Remote Sensing Perspective*. 2<sup>nd</sup> Edition. Upper Saddle River, NJ: Prentice Hall.
- Langford, M. and Unwin, D. (1994). Generating and mapping population density surfaces within a geographical information system. *Cartographic Journal* 31:21–26.
- Langford, M., Maguire, D. and Unwin, D. (1991). The areal interpolation problem: estimating population using remote sensing in a GIS framework, pp. 55-77, in: I. Masser and M. Blakemore (eds.), *Handling Geographic Information: Methodology and Potential Applications*. Harlow, Essex: Longman.
- Lillesand, T. M. and Kiefer, R. W. (1987). *Remote Sensing and Image Interpretation*. 2<sup>nd</sup> Edition. New York, NY: John Wiley and Sons.
- Longley, P.A., Goodchild, M.F., Maguire, D.J. and Rhind,, D.W. (2005). *Geographic information systems and science*. 2<sup>nd</sup> edition. New York: John Wiley & Sons.

Reibel, M. and M.E. Bufalino. (2005). A Test of Street Weighted Areal Interpolation Using Geographic Information Systems. *Environment and Planning A* 37:127-139.

Reibel, M. and Agrawal, A. (2005). Land Use Weighted Areal Interpolation. *Proceedings of the GIS Planet 2005 International Conference*, Estoril, Portugal.

Schowengerdt, R. A. (1997). *Remote Sensing: Models and Methods for Image Processing*. 2<sup>nd</sup> Edition. San Diego, CA: Academic Press.

Tobler, W. (1979). Smooth pycnophylactic interpolation for geographical regions. *Journal of the American Statistical Association* 74: 519-536.